NM FOOD/INSECT & EMERGENCY ALLERGY ACTION PLAN and MEDICATION AUTHORIZATION

Schoo	ر ا District / School Name				Dat	e			_	
Student Name			Date of Birth Studen			nt #				
*Health Care Provider Name/Title			Provider's Office Phone / FAX #					_	ice student's icture here	
Parent/Guardian			Parent's Phone #s							
Emergency Contact			Contact Phone #s							
	agnosis of Mild Allergy?	History of Asthma? No Yes (Asthma may indicate an increased risk of severe reaction) History of SEVERE Anaphylactic Reaction? No Yes, If checked YES, give epinephrine immediately! Give epinephrine if allergen was <i>likely</i> eaten, at onset of <i>any</i> symptoms or if allergen was definitely eaten even if no symptoms are noticed.								
TREATMENT PLAN	FOR ANY OF THE FOLLOWING SEVERE SYMPTOMS:LUNG:Difficulty breathing or swallowing, wheezing, coughingHEART:Dizzy, faint, confused, pale, blue, weak pulseTHROAT:Tight, hoarse, trouble breathing/swallowing, droolingMOUTH:Significant swelling of tongue, lipsSKIN:Many hives over body, widespread redness over bodyGUT:Nausea, repetitive vomiting, severe diarrhea, crampingOther:Feeling something bad is about to happen, anxiety, confusion					 FOLLOW THIS PROTOCOL: 1. INJECT EPINEPHRINE IMMEDIATELY! (Note time) 2. Call 911. Request ambulance with epinephrine. 3. Don't hang up & don't leave student 4. Give additional medications as ordered Antihistamine (if ordered below) Inhaler (Albuterol) if student has asthma 5. Lay student flat and raise legs. If breathing is difficult or vomiting, sit up or lie on their side 6. Notify School Nurse and Parent/Guardian 7. Notify Prescribing Provider / PCP 8. Student must be transported to ER 				
T	 MILD ALLERGY SYMPTOMS (IF DIAGNOSIS CONFIRMED ABOVE): MOUTH: Itchy mouth, lips, tongue and/or throat SKIN: Itchy mouth NOSE: Itchy/runny nose GUT: Mild nausea/discomfort Itchy/runny nose <l< td=""><td></td></l<>									
> THE SEVERITY OF SYMPTOMS CAN QUICKLY CHANGE. ALL SYMPTOMS OF ANAPHYLAXIS CAN POTENTIALLY PROGRESS TO A LIFE THREATENING SITUATION!!										
RDER	Epinephrine Epinephrine (0.15mg) inject intramuscularly Epinephrine (0.3mg) inject Student's weight Epi Pen Auvi Q Adrenaclick Epi Pen Auvi Q Adrenaclick Ibs. Ibs. Becond dose of epinephrine can be given 5 minutes or more after the first if symptomic Ibs. Ibs. Ibs.						Q Adrenaclic	k		
MEDICATION ORDER	Antihistamine Do not depend on antihistamines (or inhalers). When in doubt, give epinephrine and call 911. NOTE: IE NURSE IS NOT AVAILABLE. THE ABOVE TREA			Dose: Route:			INCLUDE ANXIETY, DIZZINES PALENES	SIDE EFFECTS OF EPINEPHRINE MAY INCLUDE: ANXIETY, TREMOR, PALPITATIONS, DIZZINESS, WEAKNESS, TINGLING, & PALENESS		
► NOTE: IF NURSE IS NOT AVAILABLE, THE ABOVE TREATMENT PLAN MAY BE PROVIDED BY TRAINED SCHOOL PERSONNEL FOR ANY ANAPHYLAXIS SYMPTOMS.										
MUST BE COMPLETED BY HEALTHCARE PROVIDER, PARENT, AND SCHOOL NURSE										
AUTHORIZATION	*Prescriber's Signature: Date: Date: I have complex printed Name: Phone: Phone: Complex care care care care care care care care						completed the care plan and s school personn	e reviewed this order and pleted the allergy emergency plan and shared with trained		
	administer medication(s), and contact my provider, if necessary. I assume full responsibility for providing the school with the processing description of the school staff							Signatu Medication Ex	ure / Date cpires on:	