

# HYPOGLYCAEMIA

**LOW**

Blood Glucose Level < 4.0mmol/L

**DO NOT LEAVE CHILD UNATTENDED  
DO NOT DELAY TREATMENT**

## Signs and symptoms

Note: Symptoms may not always be obvious.  
Pale, headache, shaky, sweaty, dizzy, drowsy, changes in behavior.

### Child conscious

(Able to eat hypo food)

### Child unconscious/ drowsy

(Risk of choking/unable to swallow)

### Step 1: Give fast acting carb

(as supplied or listed on management plan)

### First Aid DR<sup>S</sup>ABCD

Stay with unconscious child

### Step 2: Recheck BGL after 15 mins

If BGL < 4.0 repeat Step 1  
If BGL > 4.0 go to Step 3

**CALL AN  
AMBULANCE  
DIAL 000**

### Step 3: Give sustaining carb

(As supplied or listed on management plan)

### Contact parent/ guardian

When safe to do so

PARENT / GUARDIAN NAME \_\_\_\_\_

CONTACT No \_\_\_\_\_

## EARLY CHILDHOOD EDUCATION AND CARE SETTING

**Multiple daily injections**

Use in conjunction with management plan

# DIABETES ACTION PLAN 2017




Photo of child

CHILD'S NAME

\_\_\_\_\_  
\_\_\_\_\_

CENTRE

\_\_\_\_\_  
\_\_\_\_\_

## INSULIN

Insulin is taken 4 or more times per day.  
An injection will be needed before lunch.

**This injection requires assistance.**

Injection will be given by: \_\_\_\_\_

Injection will be given in: \_\_\_\_\_  
(Room/location)

## ROUTINE BGL CHECKING TIMES

- Anytime, anywhere in the centre
- Prior to lunch
- Anytime hypo is suspected
- Prior to planned activity

## PHYSICAL ACTIVITY

- 1 serve sustaining carb before every 30 mins of planned activity
- Vigorous activity should **not** be undertaken if BGL >15.0 and blood ketones are >1.0
- Usual play time usually doesn't require additional action, but check with parent/guardian about this

AP MD ECEC NZ

# HYPERGLYCAEMIA

**HIGH**

Blood Glucose Level >15.0mmol/L

**HIGH BGLS ARE NOT UNCOMMON**

## Signs and Symptoms

Note: Symptoms may not always be obvious.  
Some could be: increased thirst, increased urine production, poor concentration, irritability, lethargy.

### Child well

Re-check BGL in 2 hours

### Child unwell

E.g. vomiting  
Check blood ketones if able

### Encourage oral fluids, return to activity

1-2 glasses water per hour; extra toilet visits may be required.

**CONTACT PARENT/  
GUARDIAN  
TO COLLECT  
CHILD ASAP**

**In 2 hours, if BGL  
still >15.0  
call parent/guardian  
for advice**

DATE \_\_\_\_\_

HOSPITAL \_\_\_\_\_

TREATING DNS \_\_\_\_\_

CONTACT No \_\_\_\_\_